

Infrared Sensitivity in Damping Rate for Very Soft Moving Fermions in Finite Temperature QED

A. Abada^{a*} and K. Bouakaz^{b†}

^aDépartement de Physique, Ecole Normale Supérieure
BP 92 Vieux Kouba, 16050 Alger, Algeria

We calculate the fermion damping rate to second order in powers of the external momentum p in the context of QED at finite temperature using the hard-thermal-loop (HTL) summation scheme. We find that the coefficient of order p^2 is divergent in the infrared whereas the two others are finite. This result suggests that the htl-based perturbation is infrared sensitive at next-to-leading order.

At high temperature, perturbative calculations based on bare propagators and vertices lead to gauge dependent and infrared divergent results [1]. This problem has been solved within the hard thermal loop (HTL) summation technique [2,3]. The basic idea is to distinguish between momentum scales T (hard) and eT (soft), where e is the QED coupling constant. Using this method, the explicit calculation of the zero-momentum transverse gluon damping rate $\gamma_t(0)$ to next-to-leading order was performed in [4] and was shown to be finite, positive and independent of the gauge. A similar computation for the damping rate of a quark at rest was carried independently in [5] and [6].

However, recent work shows that the damping rates for very soft moving excitations in hot QCD are infrared sensitive. Indeed, an explicit calculation of the longitudinal gluon damping rate $\gamma_l(0)$ shows that this latter is different from $\gamma_t(0)$ and infrared divergent [7,8]. Also, the calculation of the transverse gluon damping rate with very soft momentum p is discussed in [9] and presented in the form $\gamma_t(p) = a_{t0} + a_{t1} \left(\frac{p}{m_g}\right)^2 + \dots$, where m_g is the soft gluon mass. The zeroth-order coefficient a_{t0} is finite and equal to $\gamma_t(0)$ found in [4], but the second-order coefficient is infrared divergent. Finally, the calculation of the damping rate for quarks to second order in the external momentum discussed in [10] suggests also that the second-order coefficient is divergent whereas the zeroth-order and the first-order ones are finite.

In this work, we calculate the damping rates of very soft fermions at next-to-leading order in finite temperature QED. We use the imaginary-time formalism with Euclidean momenta $P^\mu = (p_0, \vec{p})$ so that $P^2 = p_0^2 + p^2$ and $p = |\vec{p}|$. We follow the conventions of [11]. The fermion propagator is obtained by summing the hard-thermal-loop fermion self-energy diagrams. It is defined by ${}^*\Delta_f^{-1}(P) = \not{P} - \delta\Sigma(P)$, where $\delta\Sigma$ is the HTL in

*abada@wissal.dz

†bouakazk@caramail.com

the fermion self-energy. The effective fermion propagator can be decomposed into the two helicity eigenstates:

$${}^*\Delta_F(P) = -\frac{1}{2} \left[(\gamma^0 - i\boldsymbol{\gamma} \cdot \hat{\mathbf{p}}) \Delta_+(P) + (\gamma^0 + i\boldsymbol{\gamma} \cdot \hat{\mathbf{p}}) \Delta_-(P) \right], \quad (1)$$

where $\Delta_{\pm}(P) = \left(ip_0 \mp p - \frac{m_f^2}{2p} \left[\left(1 \mp \frac{ip_0}{p} \right) \ln \frac{ip_0 + p}{ip_0 - p} \right] \pm 2 \right)^{-1}$; $m_f = eT/\sqrt{8}$ is the fermion thermal mass. The poles of $\Delta_{\pm}(-i\omega, \mathbf{p})$ define the dispersion relation $\omega_{\pm}(p)$ for fermions to leading order in e . In Covariant gauge, the effective photon propagator is:

$${}^*\Delta_{\mu\nu}(K) = \frac{1}{G + K^2} P_{\mu\nu}^T + \frac{1}{F + K^2} P_{\mu\nu}^L + \frac{\xi}{K^2} \frac{K_{\mu} K_{\nu}}{K^2}, \quad (2)$$

where $P_{\mu\nu}^T$, $P_{\mu\nu}^L$ are the transverse and longitudinal projectors respectively and $F = \frac{2m^2 K^2}{k^2} \left(1 - \frac{ik_0}{k} Q_0 \left(\frac{ik_0}{k} \right) \right)$, $G = \frac{m^2 ik_0}{k} \left(\left(1 - \left(\frac{ik_0}{k} \right)^2 \right) Q_0 \left(\frac{ik_0}{k} \right) + \frac{ik_0}{k} \right)$, with $m = eT/\sqrt{6}$, the photon thermal mass.

The effective vertices ${}^*\Gamma$ are the sum of the tree amplitudes Γ and the hard thermal loops $\delta\Gamma$. The effective fermion-photon vertex and the effective two fermion-two photon vertex are given by:

$${}^*\Gamma^{\mu}(P_1, P_2) = \gamma^{\mu} - m_f^2 \int \frac{d\Omega_s}{4\pi} \frac{S^{\mu} \not{S}}{P_1 S P_2 S}, \quad (3)$$

$${}^*\tilde{\Gamma}^{\mu\nu}(P_1, P_2; Q) = -m_f^2 \int \frac{d\Omega_s}{4\pi} \left[\frac{1}{P_1 S} + \frac{1}{P_2 S} \right] \frac{S^{\mu} S^{\nu} \not{S}}{(P_1 + Q) S (P_2 - Q) S}. \quad (4)$$

The damping rates for fermions to second order in their very soft momentum p write:

$$\gamma_{\pm}(p) = \frac{1}{2} \left[1 \pm \frac{2}{3} \frac{p}{m_f} - \frac{2}{9} \left(\frac{p}{m_f} \right)^2 + \dots \right] \left. \text{Im } {}^*f_{\pm}(-i\omega, p) \right|_{\omega=\omega_{\pm}+i0^+}, \quad (5)$$

where ${}^*f_{\pm} = {}^*D_0 \mp {}^*D_s$ and *D_0 , *D_s are the components of the fermion effective self-energy. Therefore, the calculation of $\gamma_{\pm}(p)$ amounts to that of the imaginary part of the fermion effective self-energy.

Up to next-to-leading-order, the fermion self-energy is given in imaginary time formalism by:

$${}^*\Sigma(P) = -e^2 \text{Tr}_{\text{soft}} [{}^*\Gamma^{\mu}(P, Q) {}^*\Delta_F(Q) {}^*\Gamma^{\nu}(P, Q) {}^*\Delta_{\mu\nu}(K) + {}^*\Gamma^{\mu\nu}(P, -P; Q, -Q) {}^*\Delta_{\mu\nu}(K)], \quad (6)$$

where K is the internal soft loop momentum, $Q = P - K$ and $\text{Tr} \equiv T \sum \int \frac{d^3 k}{(2\pi)^3}$. In this latter sum, only soft values of k are allowed in the integrals. Using the expressions of the effective propagators (1, ??) and the effective vertices (3, ??), we see that there are nine contributions to ${}^*\Sigma$. Details on how to handle these is given in [10]. We find that the damping rates can be put in the following form:

$$\gamma_{\pm}(p) = -\frac{e^2 T}{8\pi} \left[a_0 \pm \frac{p}{3} a_1 + \frac{p^2}{9} a_2 + \dots \right], \quad (7)$$

with:

$$a_i = \int_{\eta}^{\infty} dk \int_{-\infty}^{+\infty} d\omega \int_{-\infty}^{+\infty} \frac{d\omega'}{\omega'} [f_i(\omega, \omega'; k) - f_{i-1, i \geq 1}(\omega, \omega'; k) \partial_{\omega} + f_{i-2, i \geq 2}(\omega, \omega'; k) [-3 \partial_{\omega} + \partial_{\omega}^2]] \delta; \quad (8)$$

$$f_0(\omega, \omega'; k) = \sum_{\epsilon=\pm} \left[-k^2 (1 - \epsilon k + \omega)^2 \rho_{\epsilon} \rho'_l + \frac{1}{2} (1 + 2\epsilon k + k^2 - \omega^2)^2 \rho_{\epsilon} \rho'_t \right] + \frac{1}{k} (k^2 - \omega^2) \rho_0 \rho'_t, \quad (9)$$

$$f_1(\omega, \omega'; k) = \sum_{\epsilon=\pm} \left[2k^2 (-1 + k^2 - 2\epsilon k \omega + \omega^2) \rho_{\epsilon} \rho'_l + \left(-\frac{2\epsilon}{k} - 3 + 2\epsilon k + 4k^2 - k^4 - (2 + 4\epsilon k + 2k^2) \omega + \left(\frac{4\epsilon}{k} + 4 + 2\epsilon k + 2k^2 \right) \omega^2 + 2\omega^3 - \left(\frac{2\epsilon}{k} + 1 \right) \omega^4 \right) \rho_{\epsilon} \rho'_t + \epsilon k^2 (1 - \epsilon k + \omega)^2 \rho_{\epsilon} \partial_k \rho'_l + \left(\frac{\epsilon}{2} + 2k + 3\epsilon k^2 + 2k^3 + \frac{\epsilon}{2} k^4 - (\epsilon + 2k + \epsilon k^2) \omega^2 + \frac{\epsilon}{2} \omega^4 \right) \rho_{\epsilon} \partial_k \rho'_t \right] - \frac{2}{k} \left(k^2 - \omega^2 + 2 \frac{\omega^3}{k^2} \right) \rho_0 \rho'_t - 2k^2 \epsilon(\omega) \delta(\omega^2 - k^2) \rho'_l + \frac{\omega}{k^2} (\omega^2 - k^2) \rho_0 \partial_k \rho'_t + 2\omega \rho_0 \partial_k \rho'_l; \quad (10)$$

$$f_2(\omega, \omega'; k) = \sum_{\epsilon=\pm} \left[\left(-\frac{9}{2} - k^2 - 6\epsilon k^3 - \frac{1}{2} k^4 - (6\epsilon k - 6k^2 + 2\epsilon k^3) \omega + (9 + k^2) \omega^2 + 6\epsilon k \omega^3 - \frac{9}{2} \omega^4 \right) \rho_{\epsilon} \rho'_l + \left(\frac{9}{2k^2} - \frac{14\epsilon}{k} - \frac{8}{3} + 4\epsilon k - \frac{19}{2} k^2 - 6\epsilon k^3 + k^4 + \left(\frac{-3}{k^2} + \frac{25\epsilon}{k} - 10 + 6\epsilon k + 9k^2 - 3\epsilon k^3 \right) \omega + \left(-\frac{6}{k^2} + \frac{2\epsilon}{k} + 23 - 6\epsilon k + k^2 \right) \omega^2 + \left(\frac{6}{k^2} - \frac{22\epsilon}{k} - 6 + 6\epsilon k \right) \omega^3 + \left(-\frac{3}{2k^2} + \frac{12\epsilon}{k} - 5 \right) \omega^4 - \left(\frac{3}{k^2} + \frac{3\epsilon}{k} \right) \omega^5 + \frac{3}{k^2} \omega^6 \right) \rho_{\epsilon} \rho'_t - k (9 - 14\epsilon k + 5k^2 + (12 - 2\epsilon k - 6k^2) \omega - (3 - 12\epsilon k) \omega^2 - 6\omega^3) \rho_{\epsilon} \partial_k \rho'_l + \left(\frac{3}{2k} + 4\epsilon + 7k + 8\epsilon k^2 + \frac{7}{2} k^3 - \left(\frac{3}{k} + 6\epsilon + 6k + 6\epsilon k^2 + 3k^3 \right) \omega - \left(\frac{3}{k} + 4\epsilon + 5k \right) \omega^2 + \left(\frac{6}{k} + 6\epsilon + 6k \right) \omega^3 + \frac{3}{2k} \omega^4 - \frac{3}{k} \omega^5 \right) \rho_{\epsilon} \partial_k \rho'_t - \frac{3}{2} k^2 (1 - \epsilon k + \omega)^2 \rho_{\epsilon} \partial_k^2 \rho'_l + \left(\frac{3}{4} + 3\epsilon k + \frac{9}{2} k^2 + 3\epsilon k^3 + \frac{3}{4} k^4 - \frac{3}{2} (1 + 2\epsilon k + k^2) \omega^2 + \frac{3}{4} \omega^4 \right) \rho_{\epsilon} \partial_k^2 \rho'_t \right] - \frac{3}{k} (k^2 - \omega^2) \rho_0 \rho'_l + \left(\frac{3}{k} + 2k + \frac{6}{k} \omega - \left(\frac{15}{k^3} + \frac{2}{k} \right) \omega^2 + \frac{18}{k^3} \omega^3 \right) \rho_0 \rho'_t + (6 - 12\omega) \rho_0 \partial_k \rho'_l + \left(-3 + 6k\omega + \frac{3}{k^2} \omega^2 - \frac{6}{k} \omega^3 \right) \rho_0 \partial_k \rho'_t + 3k \rho_0 \partial_k^2 \rho'_l - \frac{3}{2k} (k^2 - \omega^2) \rho_0 \partial_k^2 \rho'_t + 12k^2 \epsilon(\omega) \delta(\omega^2 - k^2) \rho'_l - 6|\omega| \delta(\omega^2 - k^2) \rho'_t - 6k^2 |\omega| \partial_{\omega^2} \delta(\omega^2 - k^2) \rho'_l. \quad (11)$$

In the above expression, η is an infrared cutoff of the order of the magnetic scale. $\delta = \delta(1 - \omega - \omega')$ and $\partial_{\omega} \equiv \frac{\partial}{\partial \omega}$; $\rho_{\epsilon} = \rho_{\epsilon}(k, \omega)$ and $\rho'_{t,l} = \rho_{t,l}(k, \omega')$ are the spectral densities of the effective propagators $^* \Delta_{\epsilon}$ and $^* \Delta_{t,l}$ respectively [11,12]. Also, $\rho_0 = -\frac{1}{2} \Theta(k^2 - \omega^2)$ and $\epsilon(\omega)$ is the sign function. Note that for convenience, we have set $m_f = 1$ in the above formula.

What remains to do is the numerical evaluation of a_0 , a_1 and a_2 . For this purpose, we have to perform the integrals over the frequencies ω and ω' , and then over the momentum

k . This necessitates the explicit use of the expressions of the spectral functions of the different quantities involved [11] and a delicate extraction of potential infrared divergences. All relevant details will be given elsewhere. We report here on the final result, which writes as:

$$\gamma_{\pm}(p) = \frac{e^2 T}{16\pi} \left[5.4253 \mp 0.5566 p + (-13.3136 \ln \eta + 10.4327) p^2 + \dots \right]. \quad (12)$$

We have calculated the damping rates of fermions in finite-temperature QED to next-to-leading order in HTL-summed perturbation. We have found that the coefficients of zeroth order and order p are infrared safe whereas the third one is logarithmically divergent. This result indicates that the HTL-summation scheme is perhaps not complete at this order.

REFERENCES

1. V.P. Silin, Zh. Eksp. Teor. Fiz. **38** (1960) 1577 [Sov. Phys. JETP **11** (1960) 1136]; V.N. Tsytovich, Zh. Eksp. Teor. Fiz. **40** (1961) 1775 [Sov. Phys. JETP **13** (1961) 1249]; O.K. Kalashnikov and V.V. Klimov, Yad. Fiz. **31** (1980) 1357 [Sov. J. Nucl. Phys. **31** (1980) 699]; V.V. Klimov, Yad. Fiz. **33** (1981) 1734 [Sov. J. Nucl. Phys. **33** (1981) 934]; Zh. Eksp. Teor. Fiz. **82** (1982) 336 [Sov. Phys. JETP **55** (1982) 199]; H.A. Weldon, Phys. Rev. D **26** (1982) 1394.
2. R.D. Pisarski, Phys. Rev. Lett. **63** (1989) 1129.
3. E. Braaten and R.D. Pisarski, Nucl. Phys. **B337** (1990) 569; Phys. Rev. Lett. **64** (1990) 1338; Nucl. Phys. **B339** (1990) 310.
4. E. Braaten and R.D. Pisarski, Phys. Rev. **D42** (1990) R2156.
5. R. Kobes, G. Kunstatter and K. Mak, Phys. Rev. **D45** (1992) 4632.
6. E. Braaten and R. Pisarski, Phys. Rev. **D46** (1992) 1829.
7. A. Abada, O. Azi and K. Benchallal, Phys. Lett. **B425** (1998) 158.
8. A. Abada, et O. Azi, Phys. Lett. **B** 463 (1999) 117.
9. A. Abada, O. Azi and A. Tadj, “*Damping Rate for Transverse Gluons with Finite Soft Momentum in Hot QCD*”, hep-ph/9710549, unpublished; A. Abada, O. Azi and K. Bouakaz, in writing.
10. A. Abada, K. Bouakaz, N. Daira-Aifa, Euro. Phys.J. C **18** (2001) 765-777.
11. M. Le Bellac, ‘*Thermal Field Theory*’, Cambridge University Press, 1996.
12. R. D. Pisarski, Physica **A** 158 (1989) 146; Nucl. Phys. **B309** (1988) 476.